

For the Period May 28 to June 3, 2013

Seeding continues to progress in the province and producers now have 83 per cent of the 2013 crop in the ground, right in line with the five-year (2008-2012) average of 82 per cent seeded for this time of year.

Seventy-one per cent of the crop is seeded in the southeast; 90 per cent in the southwest; 84 per cent in the east-central area; 87 per cent in west-central and northeast areas; and 89 per cent in the northwest.

Provincially, producers have seeded 89 per cent of spring wheat, 87 per cent of field peas, 86 per cent of canola, 84 per cent of durum, 81 per cent of lentils, 76 per cent of barley, 67 per cent of oats, 54 per cent of flax, 43 per cent of mustard and chickpeas and 41 per cent of canary seed.

Continuous rainfall has delayed seeding in much of the southern portion of the province with the Radville and Coronach areas receiving more than four inches of rain in the past week.

Provincially, topsoil moisture on crop land is rated as 11 per cent surplus, 79 per cent adequate and 10 per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 79 per cent adequate and 16 per cent short.

Despite most emerged crops trailing behind their normal stages of development for this time of year the majority of these crops are in good to

excellent condition. Most crop damage this week was caused by flea beetle and cutworm damage in canola crops, excess moisture and wind.

Farmers are busy trying to finish seeding, controlling weeds and moving cattle to pasture.

One year ago

Ninety-one per cent of the 2012 crop had been seeded. Producers province-wide were dealing with flea beetles and flooding issues.

Follow the 2013 Crop Report on Twitter [@SKGovAg](https://twitter.com/SKGovAg)

Seeding Progress by Crop District

CD	June 3/13	June 4/12
1a	73	90
1b	74	85
2a	70	83
2b	75	90
3ase	76	93
3asw	83	92
3an	85	92
3bs	94	94
3bn	93	96
4a	98	94
4b	88	98
5a	78	78
5b	87	76
6a	85	94
6b	82	90
7a	86	94
7b	92	98
8a	85	95
8b	86	95
9ae	94	97
9aw	90	95
9b	88	96

Seeding Progress in SK Per cent seeded All Crops

June 3/13	83
5 year avg. (2008-2012)	82
June 4/12	91
May 30/11	70
June 3/10	59
June 1/09	95
June 1/08	97
10 year avg. (2003-2012)	88

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: croreport@gov.sk.ca.

Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.



**Southeastern Saskatchewan
(Crop Districts 1, 2, and 3ASE)**

Significant rainfall in the region has further delayed seeding progress this week. Producers now have 71 per cent of the crop in the ground, up only four per cent from last week. The five-year (2008-2012) average for this time of year is 76 per cent of acres seeded.

Seventy-nine per cent of field peas have been seeded, 77 per cent of spring wheat and durum, 74 per cent of canola and lentils, 60 per cent of barley, 50 per cent of flax, 44 per cent of oats, 39 per cent of canary seed, 24 per cent of mustard and 20 per cent of chickpeas have been seeded.

Most of the earlier seeded crops have emerged and are in fair to good condition. The majority of crops are behind normal development stages for this time of year due to cool and wet field conditions. Fifty-nine per cent of fall cereals, 66 per cent of spring cereals, 71 per cent of oilseeds and 73 per cent of pulses are behind normal developmental stages for this time of year.

Rainfall in the region ranged from less than one inch to more than four inches. Continuous rainfall has damaged emerged crops and packed the soil in some areas. The Frobisher and Maryfield areas received 50 mm of rainfall, the Stoughton area 33 mm, the Moosomin area 60 mm, the Rocanville area 47 mm, the Weyburn area 79 mm, the Wilcox area 41 mm, the Moose Jaw area 28 mm, the Ceylon area 83 mm and the Radville area 102 mm. Since April 1, the Radville area has received the greatest amount of precipitation (164 mm).

Topsoil moisture conditions on cropland are rated as 34 per cent surplus, 64 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 84 per cent adequate and three per cent short. CDs 1A and 2A are reporting that 51 per cent and 73 per cent, respectively, of the cropland has surplus topsoil moisture at this time. CD 2A is also reporting that 38 per cent of the hay and pasture land has surplus topsoil moisture.

Localized flooding, high winds and frost have damaged crops in some areas this week. Some crop reporters have indicated that if more favourable weather conditions are not received in the coming days, seeding will not be completed due to wet fields in some southern areas of the region. For others in the region, seeding is wrapping up and spraying operations are beginning. Pasture growth is variable throughout the area but for the most part is slow. Fertilizer shortages continue to be a problem for those still seeding.

Farmers are busy seeding, controlling weeds and working fields.

Southeastern Saskatchewan	
Crop District	% seeded (June 3, 2013)
1A	73
1B	74
2A	70
2B	75
3ASE	76
Region average	71
five-year average (2008-2012)	76

**Southwestern Saskatchewan
(Crop Districts 3ASW, 3AN, 3B and 4)**

Seeding continues to progress in the southwest and producers now have 90 per cent of the 2013 crop seeded, up from 85 per cent last week. The five-year (2008-2012) average for this time of year is 84 per cent. Significant rainfall in much of the region has delayed further seeding progress.

Ninety per cent of lentils have been seeded, 89 per cent of durum, 85 per cent of spring wheat and canola, 84 per cent of field peas, 66 per cent of barley, 60 per cent of flax, 48 per cent of mustard, 47 per cent of chickpeas and oats, and 21 per cent of canary seed has been seeded.

Most of the earlier seeded crops have emerged and are in fair to good condition. Seventy-two per cent of fall cereals, 60 per cent of spring cereals, 55 per cent of oilseeds and 62 per cent of pulses are at normal developmental stages for this time of year.

Southern parts of the region received significant rainfall that may delay seeding completion for some producers. The Big Beaver area received 92 mm of rainfall, the Coronach area 112 mm, the Bengough area 97 mm, the Assiniboia area 57 mm, the Mortlach area 42 mm, the Coderre and Richmond areas 34 mm, the Shaunavon area 52 mm, the Admiral area 61 mm, the Rush Lake area 27 mm, the Tyner area 3 mm, the Eastend area 44 mm, the Maple Creek area 28 mm and the Gull Lake area 19 mm. The Coronach area has received the greatest amount of precipitation since April 1 at 200 mm.

Topsoil moisture for cropland is rated as nine per cent surplus, 83 per cent adequate and eight per cent short. Hay land and pasture topsoil moisture is rated as two per cent surplus, 84 per cent adequate and 14 per cent short. CD 3ASW is reporting that 30 per cent of cropland acres have surplus topsoil moisture.

Heavy rain and hail have flooded fields and access roads in some southern areas of the region. Frost and wind have damaged emerged crops in other areas, although damage is minimal for the most part. There are reports of significant damage to crops from insects such as pea leaf weevils, wireworms, cutworms and flea beetles. Northern parts of the region would welcome moisture to help replenish the topsoil. There are reports that some winter wheat fields are being re-seeded to other crops and in-crop spraying operations are beginning.

Farmers are busy seeding and controlling weeds.

Southwestern Saskatchewan	
Crop District	% Seeded (June 3, 2013)
3ASW	83
3AN	83
3BS	94
3BN	93
4A	98
4B	88
Region average	90
five-year average (2008-2012)	84

East-Central Saskatchewan (Crop Districts 5 and 6a)

Great progress was made in the east-central region this past week, thanks to warm and dry conditions. Eighty-four per cent of the crop is now in the ground, significantly up from 60 per cent last week. The five-year (2008-2012) average for this time of year is 75 per cent.

East-central Saskatchewan	
Crop District	% Seeded (June 3, 2013)
5A	78
5B	87
6A	85
Region average	84
five-year average (2008-2012)	75

Eighty-nine per cent of canola has been seeded, 88 per cent of spring wheat, 80 per cent of field peas, 78 per cent of barley, 70 per cent of flax and oats, 67 per cent of lentils, 64 per cent of durum, 37 per cent of mustard and 11 per cent of canary seed.

Most of the earlier seeded crops have emerged and are in fair to good condition. Sixty-three per cent of fall cereals, 49 per cent of spring cereals, 50 per cent of oilseeds and 52 per cent of pulses are behind normal developmental stages for this time of year.

Some areas received rainfall that may delay seeding for some producers. The Stockholm area received 38 mm of rainfall, the Langenburg area 14 mm, the Saltcoats area 20 mm, the Roblin area 15 mm, the Goodeve and Kamsack areas 12 mm, the Pelly area 20 mm, the Bethune area 1 mm and the Holdfast area 3 mm. All other areas reported no rainfall. Since April 1, the Stockholm area has received the greatest amount of precipitation (181 mm).

Cropland topsoil moisture is rated as seven per cent surplus, 78 per cent adequate and 15 per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 75 per cent adequate and 20 per cent short.

Frost and wind has caused the majority of damage this week. There have also been reports of insect damage to emerging crops from flea beetles and cutworms. Many producers will be done seeding in the next week and in-crop herbicide operations will begin soon. Some winter wheat fields are being re-seeded to other crops as winterkill damage was high. Emerged crops and pastures are advancing quickly in some areas.

Farmers are busy seeding, controlling weeds and working fields.

**West-Central Saskatchewan
(Crop Districts 6B and 7)**

Warm and dry weather conditions have allowed seeding to progress in the west-central region. Seeding is now at 87 per cent complete, up from 65 per cent last week. The five-year (2008-2012) average for this time of year is 89 per cent.

West-central Saskatchewan	
Crop District	% Seeded (June 3, 2013)
6B	82
7A	86
7B	92
Region average	87
five-year average (2008-2012)	89

Most of the region received some rainfall this week. The Kindersley area received an average of 15 mm, the Biggar and Landis areas 10 mm, the Cando and Tramping Lake areas 8 mm and the Dinsmore area 4 mm. Since April 1, the Major area has received the greatest amount of precipitation (94 mm).

Ninety-nine per cent of field peas have been seeded, 92 per cent of canola, 90 per cent of spring wheat, 80 per cent of barley, 79 per cent of durum, 72 per cent of lentils, 62 per cent of canary seed, 55 per cent of oats, 38 per cent of mustard, 32 per cent of flax, 14 per cent of chickpeas and four per cent of sunflowers.

Some of the earlier seeded crops have emerged and are in good to excellent condition; winter crops are in good condition. More than half of the crops are behind development stages for this time of year due to later seeding dates. Thirty-eight per cent of fall cereals, 62 per cent of spring cereals, 59 per cent of oilseeds and 62 per cent of pulse crops are behind normal developmental stages for this time of year.

Cropland topsoil moisture is rated as 7 per cent surplus, 83 per cent adequate and 10 per cent short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 79 per cent adequate and 15 per cent short.

No crop damage was reported to emerging crops this week. Drier conditions this week facilitated good seeding progress; however, moisture will be needed to give crops a good start. Spraying has started in many areas as pre-seeding spraying was not possible previously. Most crop reporters are indicating that seeding is finishing up for most producers in the area.

Farmers are busy finishing seeding, controlling weeds, rolling pulses, checking for flea beetles and hauling grain.

**Northeastern Saskatchewan
(Crop Districts 8 and 9AE)**

There has been little to no rainfall in the region and seeding progressed rapidly this week. Producers now have 87 per cent of the crop in the ground, up 16 per cent from last week. The five-year (2008-2012) average for this time of year is 85 per cent of acres seeded.

Northeastern Saskatchewan	
Crop District	% Seeded (June 3, 2013)
8A	85
8B	86
9AE	94
Region average	87
five-year average (2008-2012)	85

Ninety-seven per cent of spring wheat has been seeded, 84 per cent of canola, 83 per cent of field peas, 82 per cent of barley and oats, 63 per cent of flax, 49 per cent of canary seed, 36 per cent of durum and lentils, and 11 per cent of mustard.

Some of the earlier seeded crops have emerged and are in good to excellent condition; winter crops are in good to fair condition. Roughly, half of the crops are behind development stages for this time of year due to later seeding dates. Thirty-three per cent of fall cereals, 48 per cent of spring cereals, 55 per cent of oilseeds and 51 per cent of pulse crops are behind normal developmental stages for this time of year.

There was very little rainfall in the region. The Melfort area reported receiving 2 mm last Tuesday. Since April 1, the Humboldt area has received the greatest amount of precipitation (47 mm).

Topsoil moisture conditions on cropland are rated as three per cent surplus, 92 per cent adequate and five per cent short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 92 per cent adequate and seven per cent short. CDs 8A and 8B are reporting that 6 and 4 per cent, respectively, of crop land is short of topsoil moisture at this time, whereas 4 per cent per cent of crop is still experiencing surplus moisture.

Crop reporters indicate that some crops have been damaged by localized drought and insect damage by flea beetles was also reported in some areas. Seeding operations are finishing up in most areas and spraying is underway or starting across the district. Although there is still difficulty accessing areas due to excess water, there will be a need for moisture in the next week to get crops well established.

Farmers are busy seeding, controlling weeds and working fields.

**Northwestern Saskatchewan
(Crop Districts 9AW and 9B)**

Seeding progress in the northwest has quickly progressed due to warm and dry weather. Eight-nine per cent of the crop is now in the ground, which is 20 per cent more than last week. The five-year (2008-2012) average for this time of year is 91 per cent.

Northwestern Saskatchewan	
Crop District	% Seeded (June 3, 2013)
9AW	90
9B	88
Region average	89
five-year average (2008-2012)	91

Ninety-three per cent of canola has been seeded, 92 per cent of spring wheat, 91 per cent of field peas, 79 per cent of barley, 66 per cent of oats and 21 per cent of flax.

Some of the earlier seeded crops have emerged and are in good to excellent condition; winter crops range from excellent to fair condition. Fall cereals are at normal stages of development, but most spring seeded crops are behind their normal development stages for this time of year due to later seeding dates. Seventeen per cent of fall cereals, 68 per cent of spring cereals and oilseeds and 63 per cent of pulse crops are behind normal developmental stages for this time of year.


There was a small amount of precipitation received this past week all of which was in CD 9B. The Turtleford, Glaslyn and Barthel areas received five mm of rain and the Frenchman Butte area received three mm. Since April 1, the Lloydminster area has received the most precipitation for the region (48 mm).

Topsoil moisture conditions on cropland are rated as three per cent surplus, 82 per cent adequate and 15 per cent short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 66 per cent adequate and 33 per cent short. CD 9B is reporting that 39 per cent of the pasture lands is short topsoil moisture at this time.

High wind damage and flea beetle insect damage was reported for emerging crops this week, but many crops have just been seeded and have not yet emerged. Seeding is nearing the end and spraying has started in some areas. Crop reporters indicate that moisture would be welcome in the coming week to help germinate crops.


Farmers are busy seeding, spraying, hauling grain, fencing and moving cattle to pasture.

in Saskatchewan



Weekly Rainfall

for the week ending June 3, 2013



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period May 28 to June 3, 2013

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	36	95	3BN	138 A	Webb	12	67	7A	259	Snipe Lake	5	25
	3	Ennskillen	50	121		138 B	Webb	12.7	60.6		260 A	Newcombe	N/A	50
	33	Moose Creek	N/A	50		106 A	Excelsior	27	61		260 B	Newcombe	N/A	N/A
	34	Browning	23	72		106 B	Excelsior	N/A	N/A		261	Chesterfield	5	24
	61	Antler	N/A	69		167	Sask. Landing	21	56		267	St. Andrews	NIL	26
	63	Moose Mountain	47	90		168 A	Riverside	10	57		288	Pleasant Valley	1	37
	64	Brook	27	64		168 B	Riverside	7	84.4		290 A	Kindersley	5	42.5
	65	Tecumseh	33	79		226 A	Victory	8	20		290 B	Kindersley	N/A	15.5
	91	Maryfield	50	71		226 B	Victory	27.5	30		290 C	Kindersley	7	45.5
	92	Walpole	N/A	NIL		228	Lacadena	3	19		292	Milton	N/A	27
1B	122	Martin	60	95	4A	255	Coteau	NIL	33.9	7B	317 A	Marnott	N/A	N/A
	123	Silverwood	34	97		257	Monet	14.6	31.1		317 B	Marnott	NIL	53
	124	Kingsley	31	72		19	Frontier	N/A	N/A		317 C	Marnott	N/A	N/A
	125 A	Chester	32	75.5		49	White Valley	44	91		318	Mountain View	10	72
	125 B	Chester	38	87		51 A	Reno	37	97.4		320 A	Oakdale	28	69.9
	151 A	Rocanville	35	103		51 B	Reno	N/A	N/A		320 B	Oakdale	11	53
	151 B	Rocanville	47	100		79	Arlington	45	82		321	Prainedale	3	31
	153	Willowdale	N/A	N/A		109	Carmichael	33	58		347 A	Biggar	6	48
	154	Elcapo	40	101		110	Papot	23	60		347 B	Biggar	N/A	30
	155 A	Wolsley	26	81	4B	111	Maple Creek	N/A	0		350 A	Mariposa	7.6	26.4
2A	155 B	Wolsley	12	67		139	Gull Lake	19	67		350 B	Mariposa	1	30
	67	Weyburn	73	134		141	Big Stick	29	67		351	Progress	2	32.5
	68	Brokenshell	84	149		142	Enterprise	34	91		352	Heart's Hill	4	94
	97	Wellington	24.5	63.5		160	Pithville	N/A	50		377	Glenside	2	45
	99	Caledonia	N/A	N/A		231	Happyday	7	44		378 A	Rosemount	5	43
	127 A	Francis	17	51.5		183 A	Fertile Belt	38	180.5		378 B	Rosemount	8	46
	127 B	Francis	7.5	28.5		183 B	Fertile Belt	N/A	68		379	Reford	10	28
	129	Bratt's Lake	41	76		186	Abermethy	5	44		381	Grass Lake	N/A	5.1
	131 A	Baldon	26	84		211	Churchbridge	14	58		382	Eye Hill	1.4	16
	131 B	Baldon	30	93	5A	213	Salkcoats	20	81		409 A	Buffalo	4	33
2B	156 A	Indian Head	14	31		214	Cana	N/A	25		409 B	Buffalo	N/A	N/A
	156 B	Indian Head	12	79		216	Tullymet	N/A	48.3	8A	410	Round Valley	NIL	8
	160 A	Pense	N/A	N/A		217 A	Lipton	N/A	N/A		411	Senlac	N/A	N/A
	160 B	Pense	10	69.5		241	Calder	15	67		395	Porcupine	NIL	26
	161	Moose Jaw	N/A	16		243	Wallace	N/A	21		397	Barner Valley	NIL	37.4
	162	Caron	16	63		245 A	Garry	N/A	30		428	Star City	NIL	21
	191	Marquis	5	29		245 B	Garry	14	58		456	Arboreld	NIL	16
	38 A	Launer	116	164		245 C	Garry	2	34		457	Connaught	N/A	26
	38 B	Launer	82	137		246	Buna Bon Accord	NIL	30		458	Willow Creek	NIL	5
	39 A	The Gap	83	123		247	Kelross	3	28		486	Moose Range	NIL	19
	39 B	The Gap	92	146	5B	248	Touchwood	N/A	17		487 A	Nipawin	NIL	14
3ASE	10	Happy Valley	92	117		271	Coke	12	55		487 B	Nipawin	N/A	N/A
	11	Hart Butte	111.8	200.3		273	Sliding Hills	N/A	36.5	8B	369	St. Peter	NIL	14
	12	Poplar Valley	82	145		277	Emerald	1	27		370	Humboldt	NIL	47
	40 A	Bengough	95	154		305	Invermay	NIL	25.5		371	Bayne	NIL	20
	40 B	Bengough	98	143		307	Elitros	NIL	41		372	Grant	NIL	36.5
	42	Willow Bunch	61	127		308 A	Big Quill	NIL	18		400 A	Three Lakes	N/A	0
	43 A	Old Post	56	126		308 B	Big Quill	NIL	32		400 B	Three Lakes	NIL	34
	43 B	Old Post	1	N/A		331	Livingston	20	87		402	Fish Creek	NIL	30
	70	Key West	N/A	15.3		333	Clayton	N/A	39		429	Flett's Springs	2	15
	71	Escol	57	57		334	Freezeville	2	35.2		431	St. Louis	N/A	N/A
	72 A	Lake of the Rivers	N/A	N/A		336	Sasman	NIL	30	9AE	459	Kinistino	N/A	6
3ASW	72 B	Lake of the Rivers	N/A	NIL		337 A	Lakeview	NIL	42.5		480	Birch Hills	NIL	18.4
	73 A	Stonehenge	55.9	119.6		337 B	Lakeview	N/A	9.5		481	Prince Albert	N/A	22.4
	73 B	Stonehenge	47	124		338	Lakeside	NIL	40		488	Torch River	NIL	2
	74 A	Wood River	62	107.5		366	Kelvington	NIL	46		491	Buckland	N/A	40
	74 B	Wood River	N/A	3		367	Ponass Lake	NIL	52.5	9AW	406	Mayfield	NIL	34
	74 C	Wood River	N/A	N/A		190 A	Duffern	1	45		435	Redberry	NIL	32
	101	Terrill	60	103	6A	190 B	Duffern	NIL	61.5		436	Douglas	NIL	44
	102	Lake Johnston	18	42.8		190 C	Duffern	NIL	46		463	Duck Lake	NIL	29
	103 A	Sutton	24	55		190 D	Duffern	N/A	42		466	Meeting Lake	N/A	N/A
	103 B	Sutton	N/A	N/A		219	Longlakeston	NIL	17		467 A	Round Hill	NIL	35
	132 A	Hillsborough	42	70		220	Mickillop	NIL	50		467 B	Round Hill	NIL	17
	132 B	Hillsborough	25	49.3		221	Sarnia	2.6	34.8		494	Canwood	N/A	N/A
	134	Shamrock	34	104		222	Craik	NIL	53		496 A	Spinwood	N/A	N/A
	164	Chaplin	17	17		251	Big Arm	NIL	37		496 B	Spinwood	N/A	35.9
	193 A	Eyebrow	NIL	24		252	Arm River	NIL	41	9B	497	Medstead	N/A	8
	193 B	Eyebrow	NIL	38.2		279	Mount Hope	NIL	26		438	Battle River	NIL	23
3BS	194	Enfield	N/A	N/A		282	McCraney	NIL	45		440	Hillsdale	N/A	19
	224	Maple Bush	1.5	36.5		309	Praine Rose	NIL	48		442	Manitou Lake	4	41
	17 A	Val Marie	44	140		310	Ustome	N/A	N/A		498 A	Parkdale	5	19.5
	17 B	Val Marie	N/A	N/A		313	Lost River	N/A	N/A		498 B	Parkdale	NIL	17
	45	Mankota	74	124.5		339	Leroy	NIL	52.8		499 A	Mervin	5	25.4
	75 A	Pinto Creek	38	72		340	Wolverme	NIL	77		499 B	Mervin	2	23.5
	75 B	Pinto Creek	N/A	22		341	Viscount	N/A	20		501 A	Frenchman Butte	2	47
	76	Auvergne	30	71		343 A	Blucher	NIL	69.5		501 B	Frenchman Butte	9	32
	77	Wise Creek	61	133		343 B	Blucher	N/A	0		501 C	Frenchman Butte	3	43
	78 A	Grassy Creek	43	84	6B	283	Rosedale	NIL	50.5		502 A	Britannia	N/A	N/A
3BS	78 B	Grassy Creek	60	96		284	Rudy	NIL	39		502 B	Britannia	2	48
	105	Glenbairn	51	84		285	Fertile Valley	NIL	33		561	Loon Lake	5	29
	106	Whiska Creek	38	68		286	Milden	4	17		588 A	Meadow Lake	N/A	29
	107 A	Lac Pelletier	N/A	37		314	Dundum	1	71		588 B	Meadow Lake	NIL	38
	107 B	Lac Pelletier	N/A	43.5		344	Corman Park	NIL	22		588 C	Meadow Lake	NIL	34
	108	Bone Creek	35	76		346	Perdue	NIL	34		588 D	Meadow Lake	N/A	31
						376 A	Eagle Creek	N/A	35		588 E	Meadow Lake	N/A	N/A
						376 B	Eagle Creek	NIL	47		622	Beaver River	NIL	43
						403 A	Routhern	NIL	12					
						403 B	Routhern	NIL	39					


Municipality No: A, B, C, D and E - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2013


To: June 3, 2013



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions


June 4, 2013



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions

June 4, 2013



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.